

Chromatic Ramsey number of acyclic hypergraphs

András Gyárfás*

Alfréd Rényi Institute of Mathematics
Hungarian Academy of Sciences
Budapest, P.O. Box 127
Budapest, Hungary, H-1364
gyarf@renyi.hu

Alexander W. N. Riasanovsky
University of Pennsylvania
Department of Mathematics
Philadelphia, PA 19104, USA
alexneal@math.upenn.edu

Melissa U. Sherman-Bennett
Bard College at Simon's Rock
Department of Mathematics
Great Barrington, MA 01230, USA
mshermanbennett12@simons-rock.edu

June 16, 2015

*Advisor of the research conducted in the Research Opportunities Program of Budapest Semesters in Mathematics, Spring 2015.

Abstract

Suppose that T is an acyclic r -uniform hypergraph, with $r \geq 2$. We define the (t -color) chromatic Ramsey number $\chi(T, t)$ as the smallest m with the following property: if the edges of any m -chromatic r -uniform hypergraph are colored with t colors in any manner, there is a monochromatic copy of T . We observe that $\chi(T, t)$ is well defined and

$$\left\lceil \frac{R^r(T, t) - 1}{r - 1} \right\rceil + 1 \leq \chi(T, t) \leq |E(T)|^t + 1$$

where $R^r(T, t)$ is the t -color Ramsey number of H . We give linear upper bounds for $\chi(T, t)$ when T is a matching or star, proving that for $r \geq 2, k \geq 1, t \geq 1$, $\chi(M_k^r, t) \leq (t - 1)(k - 1) + 2k$ and $\chi(S_k^r, t) \leq t(k - 1) + 2$ where M_k^r and S_k^r are, respectively, the r -uniform matching and star with k edges.

The general bounds are improved for 3-uniform hypergraphs. We prove that $\chi(M_k^3, 2) = 2k$, extending a special case of Alon-Frankl-Lovász' theorem. We also prove that $\chi(S_2^3, t) \leq t + 1$, which is sharp for $t = 2, 3$. This is a corollary of a more general result. We define $H^{[1]}$ as the 1-intersection graph of H , whose vertices represent hyperedges and whose edges represent intersections of hyperedges in exactly one vertex. We prove that $\chi(H) \leq \chi(H^{[1]})$ for any 3-uniform hypergraph H (assuming $\chi(H^{[1]}) \geq 2$). The proof uses the list coloring version of Brooks' theorem.

1 Introduction

A hypergraph $H = (V, E)$ is a set V of *vertices* together with a nonempty set E of subsets of V , which are called *edges*. In this paper, we will assume that for each $e \in E$, $|e| \geq 2$. If $|e| = r$ for each $e \in E$, then H is *r -uniform*; a 2-uniform H is a graph. A hypergraph H is *acyclic* if H contains no cycles (including 2-cycles which are two edges intersecting in at least two vertices). If H is a connected acyclic hypergraph, we say that H is a *tree*. In particular, a *star* is a tree in which one vertex is common to every edge. A *matching* is a hypergraph consisting of pairwise disjoint edges, with every vertex belonging to some edge. We denote by S_k^r and M_k^r the r -uniform k -edge star and matching, respectively.

For a positive integer k , a function $c : V \rightarrow \{1, \dots, k\}$ is called a k -coloring of H . A coloring c is *proper* if no edge of H is monochromatic under c . The chromatic number of H , denoted $\chi(H)$, is the least $m \geq 1$ for which there exists a proper m -coloring of H and in this case, we say that H is m -chromatic. Given $H = (V, E)$, a partition $\{E_1, \dots, E_t\}$ of E into t parts is called a t -edge-coloring of H . For r -uniform hypergraphs H_1, H_2, \dots, H_t , the (t -color) Ramsey number $R^r(H_1, H_2, \dots, H_t)$ is the smallest integer n for which the following is true: under any t -edge-coloring of the complete r -uniform hypergraph K_n^r , there is a monochromatic copy of H_i in color i for some $i \in \{1, 2, \dots, t\}$. When all $H_i = H$ we use the notation $R^r(H, t)$.

Bialostocki and the senior author of this paper extended two well-known results in Ramsey theory from complete host graph K_n to arbitrary n -chromatic

graphs [4]. One extends a remark of Erdős and Rado stating that in any 2-coloring of the edges of a complete graph K_n there is a monochromatic spanning tree. The other is the extension of the result of Cockayne and Lorimer [5] about the t -color Ramsey number of matchings. In [8], an acyclic graph H is defined as t -good if every t -edge coloring of any $R^2(H, t)$ -chromatic graph contains a monochromatic copy of H . Matchings are t -good for every t [4] and in [8] it was proved that stars are t -good, as well as the path P_4 (except possibly for $t = 3$). Additionally, P_5, P_6, P_7 are 2-good. In fact, as remarked in [4], there is no known example of an acyclic H that is not t -good.

In this paper, we explore a similar extension of Ramsey theory for hypergraphs, motivating the following definition.

Definition 1. *Suppose that T is an acyclic r -uniform hypergraph. Let $\chi(T, t)$ be the smallest m with the following property: under any t -edge-coloring of any m -chromatic r -uniform hypergraph, there is a monochromatic copy of T .*

We call $\chi(T, t)$ the *chromatic Ramsey number of T* . It follows from the existence of hypergraphs of large girth and chromatic number that the chromatic Ramsey number can be defined only for acyclic hypergraphs.

2 New results

First we note that $\chi(T, t)$ is well-defined for any r -uniform tree T and any $t \geq 1$, as an upper bound comes easily from the following result.

Lemma A. ([10],[12]) *If H is r -uniform with $\chi(H) \geq k + 1$, then H contains a copy of any r -uniform tree on k edges.*

Theorem 2. *For any r -uniform tree T , $\chi(T, t) \leq |E(T)|^t + 1$.*

Proof. Fix $t \geq 1$. Let T be an r -uniform tree with k edges and let $H = (V, E)$ be a hypergraph with $\chi(H) \geq k^t + 1$. Let $E = E_1 \dot{\cup} \dots \dot{\cup} E_t$ be a t -coloring of its edges.

Then $\chi((V, E_1)) \cdots \chi((V, E_t)) \geq k^t + 1$ and without loss of generality, $\chi((V, E_1)) \geq k + 1$. Then by Lemma A, (V, E_1) contains a copy of T . \square

Since any r -uniform acyclic hypergraph T may be found in some r -uniform tree T' and $\chi(T, t) \leq \chi(T', t)$, $\chi(T, t)$ is in fact well-defined for any r -uniform acyclic hypergraph and for any $t \geq 1$. Observe the following natural lower bound of $\chi(T, t)$. Let $L(T, t, r) := \left\lceil \frac{R^r(T, t) - 1}{r - 1} \right\rceil + 1$.

Proposition 3. $L(T, t, r) \leq \chi(T, t)$

Proof. Let $N := R^r(T, t) - 1$. By the definition of the Ramsey number, there is a t -coloring of the edges of K_N^r without a monochromatic T . Since $\chi(K_N^r) = \left\lceil \frac{N}{r-1} \right\rceil$, the proposition follows. \square

The notion of t -good graphs can be naturally extended to hypergraphs using Proposition 3. An acyclic r -uniform hypergraph T is called t -good if every t -edge coloring of any $L(T, t, r)$ -chromatic r -uniform hypergraph contains a monochromatic copy of T . In other words, T is t -good if $L(T, t, r) = \chi(T, t)$. Note that for $r = 2$, this gives the definition of good graphs. Although it is unlikely that all acyclic hypergraphs are t -good, we have no counterexamples.

For special families of r -uniform acyclic hypergraphs, we found linear upper bounds for $\chi(T, t)$, improving upon the general exponential upper bound above. Surprisingly, most of the bounds attained do not depend on r .

2.1 Matchings

Indispensable in this section is the following well-known result of Alon, Frankl, and Lovász (originally conjectured by Erdős).

Theorem B. ([1]) For $r \geq 2, k \geq 1, t \geq 1$,

$$R^r(M_k^r, t) = (t-1)(k-1) + kr.$$

Note that special cases of Theorem B include $r = 2$ [5], $k = 2$ [13], $t = 2$ [2], [9].

We obtain the following linear upper bound for matchings using Theorem B.

Theorem 4. For $r \geq 2, k \geq 1, t \geq 1$, $\chi(M_k^r, t) \leq (t-1)(k-1) + 2k$. Equality holds for $r = 2$.

Proof. Let $H = (V, E)$ be an r -uniform hypergraph with $\chi(H) = p$ where $p := R^r(M_k^2, t)$. By Theorem B, $p = (t-1)(k-1) + 2k$. Consider any t -edge coloring $\{E_1, \dots, E_t\}$ of H and any proper coloring c of H obtained by the greedy algorithm (under any ordering of its vertices). Clearly c uses at least p colors and for any $1 \leq i < j \leq p$ there is an edge e_{ij} in H whose vertices are colored with color i apart from a single vertex which is colored with j . Let $\{F_1, \dots, F_t\}$ be a t -edge-coloring of K_p^2 defined so that $F_s := \{\{i, j\}, 1 \leq i < j \leq p, e_{ij} \in E_s\}$ for each s , $1 \leq s \leq t$. From the definition of p , Theorem B (in fact the Cockayne-Lorimer Theorem suffices) implies that there is a monochromatic M_k^2 in K_p . Observe that

$$\{e_{ij} : \{i, j\} \in M_k^2\}$$

is a set of k pairwise disjoint edges in H in the same partition class of $\{E_1, \dots, E_t\}$. This completes the proof that $\chi(M_k^r, t) \leq (t-1)(k-1) + 2k$. The lower bound $R^2(M_k^2, t) \leq \chi(M_k^2, t)$ implies equality in the $r = 2$ case. \square

Next we tighten this bound, provided $r \geq 3$ and $t = 2$.

Theorem 5. For $r \geq 3$ and $k \geq 1$, $\chi(M_k^r, 2) \leq 2k$.

Proof. We fix $r \geq 3$ and proceed by induction on k . Suppose $k = 1$ and let H be some r -uniform hypergraph with $\chi(H) \geq 2$. Then any 2-edge-coloring of H contains a single monochromatic edge since H has at least one edge. Now

suppose the theorem is true for $k - 1 \geq 1$ and let $H = (V, E)$ be r -uniform with $\chi(H) \geq 2k$. Without loss of generality, H is connected. Fix some 2-edge-coloring $\{E_1, E_2\}$ of H , calling the edges of E_1 “red” and the edges of E_2 “blue”. If E_1 or E_2 is empty, then Theorem 4 with $t = 1$ implies the desired bound.

So we may assume otherwise, and there exist edges $e, f \in E$ with e red and f blue. Let $s := |e \cap f|$ and $A := e \cup f$. If $H[A]$ is 2-colorable, then $\chi(H - A) \geq \chi(H) - 2 = 2(k - 1)$ so by induction we find a monochromatic M_{k-1}^r matching in $H - A$. Without loss of generality, M_{k-1}^r is red and $M_{k-1}^r + e$ is a red M_k^r in H .

If $s > 1$, then $|A| = 2r - s \leq 2r - 2$ thus $H[A]$ is certainly 2-colorable and the induction works. If $s = 1$ and $H[A]$ is not 2-colorable then $H[A]$ is K_{2r-1}^r . Writing $e = \{w, u_1, \dots, u_{r-1}\}$ and $f = \{w, v_1, \dots, v_{r-1}\}$, the edge $g = \{w\} \cup \{u_1, u_3, \dots\} \cup \{v_2, v_4, \dots\} \in E(H)$. Without loss of generality, g is red and $|g \cap f| = 1 + \lfloor (r - 1)/2 \rfloor \geq 2$ since $r \geq 3$. So the previous case applies to the red edge g and blue edge f . Finally, if $s = 0$ and $H[A]$ is not 2-colorable there must be $g \in H[A]$ that intersects both e and f . Then either e, g or f, g is a pair of edges of different color that intersect, and a previous case can be applied again. \square

Corollary 6. $\chi(M_k^3, 2) = 2k$.

Proof. The upper bound is given by Theorem 5. The lower bound comes from Proposition 3 and Theorem B:

$$L(M_k^3, 2, 3) = \left\lceil \frac{k - 1 + 3k - 1}{2} \right\rceil + 1 = 2k \leq \chi(M_k^3, 2).$$

\square

Corollary 7. For $r \geq 3$, $\chi(M_2^r, 2) = 4$.

Proof. As in Corollary 6, the upper bound comes from Theorem 5 and the lower bound from

$$L(M_2^r, 2, r) = \left\lceil \frac{2r - 1}{r - 1} \right\rceil + 1 = 4.$$

\square

It is worth noting that Corollary 7 does not extend Theorem B to the chromatic Ramsey number setting for $r \geq 4$. Indeed, for $r = 4$, the lower bound $\lceil \frac{2r}{r-1} \rceil + 1$ of Proposition 3 is 4 and the bound $\lceil \frac{1+2r}{r-1} \rceil$ derived from Theorem B is 3.

2.2 Stars

Theorem 8. For $r \geq 2, k \geq 1, t \geq 1$, $\chi(S_k^r, t) \leq t(k - 1) + 2$.

Proof. Fix $t, k \geq 1$ and let $p := t(k-1) + 2$. Suppose that H is r -uniform with $\chi(H) \geq p$ and its edges are t -colored. By Lemma A, $\chi(S_{p-1}^r, 1) \leq p$, so we can find a copy of S_{p-1}^r in H . By the pigeonhole principle, k of the edges of S_{p-1}^r have the same color, and together they are a monochromatic copy of S_k^r . \square

How good is the estimate of Theorem 8? Notice first that for $t = 1$ it is sharp.

Proposition 9. $\chi(S_k^r, 1) = k + 1$

Proof. Consider the complete hypergraph $K = K_{k(r-1)}^r$. Clearly, $\chi(K) = k$ and S_k^r is not a subgraph of K , as its vertex set is too large. \square

If $t = 2$, Theorem 8 gives $\chi(S_k^r, 2) \leq 2k$. For $r = 2$ and odd k , this is a sharp estimate. For $k = 1$, this is trivial; for $k \geq 3$, the complete graph K_{2k-1}^2 can be partitioned into $2(k-1)$ -regular subgraphs. However, for even $k \geq 2$, $\chi(S_k^2, 2) = 2k - 1$.

An interesting problem arises when $T = S_2^r$ with $r \geq 3$, as Theorem 8 gives the relatively low upper bound 4. Can we decrease this bound? Namely:

Question 10. *Is $\chi(S_2^r, 2) = 3$?*

For $r = 3$ the positive answer (Corollary 14) comes from a more general result, Theorem 13 below. We first need a definition.

Definition 11. *Let $H = (V(H), E(H))$ be a hypergraph. The 1-intersection graph of H is denoted $H^{[1]}$, where $V(H^{[1]}) = E(H)$ and*

$$E(H^{[1]}) = \{(e, f) : e, f \in E(H) \text{ and } |e \cap f| = 1\}.$$

It is well-known that if $H^{[1]}$ is trivial, i.e., no two edges of H intersect in exactly one vertex, then H is 2-colorable ([14], Exercise 13.33). Note that the stronger statement $\chi(H) \leq \chi(H^{[1]}) + 1$ follows from applying the greedy coloring algorithm in any order of the vertices of H .

Question 12. *Let $r \geq 3$. Is it true that $\chi(H) \leq \chi(H^{[1]})$ for any r -uniform hypergraph H , provided $H^{[1]}$ is nontrivial?*

Our main result is the positive answer to Question 12 for the 3-uniform case.

Theorem 13. *If H is a 3-uniform hypergraph with $\chi(H^{[1]}) \geq 2$ then $\chi(H) \leq \chi(H^{[1]})$.*

Corollary 14. *For $t \geq 1$, $\chi(S_2^3, t) \leq t + 1$.*

The case $t = 2$ of Corollary 14 was the initial aim of the research in this paper and it was proved first by Zoltán Füredi [7]. Our proof of Theorem 13 uses his observation (Lemma 15 below) and the list-coloring version of Brooks' theorem. Corollary 14 is obviously sharp for $t = 2$; it follows from Proposition 3 that it is also sharp for $t = 3$, because $R^3(S_2^3, 3) = 6$ ([3]). It would be interesting to see whether Corollary 14 is true for any S_2^r (in particular for $r = 4, t = 2$) as this is equivalent to the statement that r -uniform hypergraphs with bipartite 1-intersection graphs are 2-colorable.

3 Proof of Theorem 13

In this section, we use the phrase “triple system” for a 3-uniform hypergraph. The word “triple” will take the place of “edge” so that “edge” may be reserved for graphs. Our goal is to construct a proper t -coloring of H from a proper t -coloring of $H^{[1]}$. Note that a partition of $E(H)$ into classes E_1, E_2, \dots, E_t such that for any i , $1 \leq i \leq t$, no two edges of E_i 1-intersect is precisely a proper t -coloring of $H^{[1]}$. Let B_k denote the triple system with k edges intersecting pairwise in the vertices $\{v, w\}$, called the *base* of B_k . A B -component (also, B_k -component) is a triple system which is isomorphic to B_k for some $k \geq 1$. A K -component is either three or four distinct triples on four vertices. A triple system is connected if for every partition of its vertices into two nonempty parts, there is a triple intersecting both parts. Every triple system can be uniquely decomposed into pairwise disjoint connected parts, called components. Components with one vertex are called trivial components.

Lemma 15. *Let C be a nontrivial component in a triple system without 1-intersections. Then C is either a B -component or a K -component.*

Proof. If C has at most four vertices then $1 \leq |E(C)| \leq 4$ (where $E(C)$ is here considered as a set, not a multiset) and by inspection, C is either B_1, B_2 , or a K -component. Assume C has at least five vertices and select the maximum m such that $e_1, e_2, \dots, e_m \in E(C)$ are distinct triples intersecting in a two-element set, say in $\{x, y\}$. Clearly, $m \geq 2$. Then $A = \cup_{i=1}^m e_i$ must cover all vertices of C , as otherwise there is an uncovered vertex z and a triple f containing z and intersecting A , since C is a component. However, from $m \geq 2$ and the intersection condition, $f \cap A = \{x, y\}$ follows, contradicting the choice of m . Thus $A = V(C)$ and from $|V(C)| \geq 5$ we have $m \geq 3$. It is obvious that any triple of C different from the e_i 's would intersect some e_i in one vertex, violating the intersection condition. Thus C is isomorphic to B_m , concluding the proof. \square

A multigraph G is called a *skeleton* of a triple system H if every triple contains at least one edge of G . We may assume that $V(H) = V(G)$. A *matching* in a multigraph is a set of pairwise disjoint edges. A *factorized complete graph* is a complete graph on $2m$ vertices whose edge set is partitioned into $2m - 1$ matchings. The following lemma allows us to define a special skeleton of triple systems.

Lemma 16. *Suppose that H is a triple system with $\chi(H^{[1]}) = t \geq 2$ and let H_1, H_2, \dots, H_t be a partition of H into triple systems without 1-intersections. There exists a skeleton G of H with the following properties.*

1. $E(G) = \cup_{i=1}^t M_i$ where each M_i is a matching and a skeleton of H_i .
2. For $1 \leq i \leq t$, edges of M_i are the bases of all B -components of H_i and two disjoint vertex pairs from all K -components of H_i .

3. If $K^* = K_{t+1} \subset G$ then K^* is a connected component of G factorized by the M_i 's and there is $e \in M_1 \cap E(K^*)$ such that e is from a B -component of H_1 .

Proof. From Lemma 15 we can define M_i by selecting the base edges from every B -component of H_i and selecting two disjoint pairs from every K -component of H_i . The resulting multigraph is clearly a skeleton of H and satisfies properties 1 and 2. We will select the disjoint pairs from the K -components so that property 3 also holds. Notice that $K^* = K_{t+1} \subset G$ must form a connected component in G because it is a t -regular subgraph of a graph of maximum degree t . Also, K_{t+1} is factorized by the M_i 's because the union of t matchings can cover at most $\frac{t(t+1)}{2} = \binom{t+1}{2}$ edges of K_{t+1} , therefore every edge of K_{t+1} must be covered exactly once by the M_i 's. Thus we have to ensure only that there is $e \in M_1 \cap E(K^*)$ with e from a B -component of H_1 . For convenience, we say that a $K^* = K_{t+1}$ is a *bad component* if such e does not exist.

Select a skeleton S as described in the previous paragraph such that p , the number of bad components, is as small as possible. Suppose that $(x, y) \in M_1$ is in a bad component U . In other words, (x, y) is in a K -component of H_1 , where $V(K) = \{x, y, u, v\}$ and $(u, v) \in M_1$. Now we replace these two pairs by the pairs $(x, u), (y, v)$ to form a new M_1 . After this switch, U is no longer a bad component. In fact, either U becomes a new component on the same vertex set (if (u, v) was in U) or U melds with another component into a new component. In both cases, no new bad components are created and in the new skeleton there are fewer than p bad components. This contradiction shows that $p = 0$ and proves the lemma. \square

Proof of Theorem 13. Let H be a triple system with $t := \chi(H^{[1]}) \geq 2$ and partition H into H_1, \dots, H_t so that each H_i is without 1-intersections. Let G be a skeleton of H with the properties ensured by Lemma 16.

Let G' be a connected component of G . By Brooks' Theorem, if G' is not the complete graph K_{t+1} or an odd cycle (if $t = 2$), $\chi(G') \leq \Delta(G') \leq t$.

Suppose first that t is even. Now $G' \neq K_{t+1}$ because that would contradict property 3 in Lemma 16: K_{t+1} cannot be factorized into matchings. Also, for $t = 2$, G' cannot be an odd cycle since odd cycles are not the union of two matchings. Thus every connected component of G is at most t -chromatic, therefore $\chi(G) \leq t$. Since G is a skeleton of H , this implies $\chi(H) \leq t$, concluding the proof for the case when t is even.

Suppose that t is odd, $t \geq 3$. In this case the previous argument does not work when some connected component $G' = K_{t+1} \subset G$. However, from Lemma 16, every K_{t+1} -component C_i of G has an edge $(x_i, y_i) \in M_1$ that is the base of a B -component in H_1 . Define the vertex coloring c on $X = \cup_{i=1}^m V(C_i)$ by $c(x_i) = c(y_i) = 1$ and by coloring all the other vertices of all C_i 's with $2, \dots, t$.

Let F be the subgraph of G spanned by $V(G) \setminus X$ and define

$$Z := \{z \in V(F) : \{x_i, y_i, z\} \in E(H_1) \text{ for some } 1 \leq i \leq m\}.$$

Since for every $z \in Z$ there is a triple $T = (x_i, y_i, z) \in H_1$ in a B -component of H_1 (with base (x_i, y_i)), $(z, u) \in M_1$ is impossible for any $u \in V(G)$, since otherwise T and the triple of H_1 containing (z, u) would 1-intersect in z . Thus $d_G(v) \leq t - 1$ for $z \in Z$. Also, $d_G(v) \leq t$ for all $v \in V(F) \setminus Z$.

We claim that with lists $L(z) := \{2, \dots, t\}$ for $z \in Z$ and $L(v) := \{1, \dots, t\}$ for $v \in V(F) \setminus Z$, F is L -choosable. We use the reduction argument present in many coloring proofs (see, for example, the very recent survey paper [6]).

Suppose F is not L -choosable and let F' be a minimal induced subgraph of F which fails to be L -choosable. We may assume that any $z \in V(F') \cap Z$ has $d_{F'}(z) = t - 1$ (otherwise we may L -choose $F' - z$, add z back and properly color it). Likewise we may assume $d_{F'}(v) = t$ for all $v \in V(F') \setminus Z$. By the degree-choosability version of Brooks' theorem (see [11], Lemma 1 or [6], Theorem 11), F' is a Gallai tree: a graph whose blocks are complete graphs or odd cycles.

Let A be a block of F' . Then $A \neq K_{t+1}$ because all K_{t+1} -components of G are in X . Since all vertex degrees in F' are t or $t - 1$, A is either an odd cycle (if $t = 3$) or A is a K_t . A must contain an edge $e \in M_1$. Otherwise M_2, \dots, M_t would cover the edges of A , a contradiction in either case. If A is an endblock then by the degree requirements, either

$$V(A) \cap (V(F) \setminus Z) = \{w\}$$

where w is the unique cut point of A or $V(A) \subset Z$. In both cases an endpoint of e must be in Z . Then there exists some triple $\{x_i, y_i, z\} \in H_1$ which 1-intersects with the triple of H_1 containing e , a contradiction, proving that F is L -choosable.

Let $c' : V(F) \rightarrow \{1, \dots, t\}$ be an L -coloring of F . We extend c from X to $V(H)$ by setting $c(v) := c'(v)$ for all $v \in V(F)$. Observe that c properly colors all edges of G except for the edges of the form (x_i, y_i) which are monochromatic in color 1. Since G is a skeleton, every triple of H is properly colored except possibly the triples in the form (x_i, y_i, x) .

We claim that $c(x) \neq 1$. Suppose to the contrary that $c(x) = 1$. If $x \in X$ then $x \in \{x_j, y_j\}$ for some $j \neq i$, but this is impossible because the bases $(x_i, y_i), (x_j, y_j)$ are from different B -components of H_1 . If $x \notin X$ then $x \in Z$ from the definition of Z . However, $1 \notin L(x)$ for $x \in Z$ and this proves the claim.

Therefore c is a proper t -coloring of H and this completes the proof. \square

References

- [1] N. Alon, P. Frankl, L. Lovász, The chromatic number of Kneser hypergraphs, *Transactions of the American Mathematical Society* 298 (1986) 359-370.
- [2] N. Alon, P. Frankl, Families in which disjoint sets have large union, in Combinatorial Mathematics; Proceedings of the Third International Conference, New York, 1985 *Ann. New York Acad. Sci.* 555 (1989) 9-16.
- [3] M. Axenovich, A. Gyárfás, H. Liu, D. Mubayi, Multicolor Ramsey numbers for triple systems, *Discrete Mathematics* **322** (2014) 133-136.
- [4] A. Bialostocki, A. Gyárfás, Replacing the host K_n by n -chromatic graphs in Ramsey type results, *arXiv:1506.04495v1*.
- [5] E. J. Cockayne, P. J. Lorimer, The Ramsey number for stripes, *Journal of Australian Mathematical Society, series A.* **19** (1975) 252-256.
- [6] D. W. Cranston, L. Rabern, Brooks' theorem and beyond, *arXiv:1403.0479v1*.
- [7] Z. Füredi, personal communication.
- [8] P. Garrison, Good graph hunting, submitted.
- [9] A. Gyárfás, On the Ramsey number of disjoint hyperedges, *MTA SZTAKI Working Paper* August, 1983.
- [10] A. Gyárfás, J. Lehel, Trees in greedy coloring of hypergraphs, *Discrete Mathematics* **311** (2011) 208-209.
- [11] A. V. Kostochka, M. Stiebitz, B. Wirth, The colour theorems of Brooks and Gallai extended, *Discrete Mathematics* **162** (1996) 299-303.
- [12] Po-Shen Loh, A note on embedding hypertrees, *Electronic Journal of Combinatorics* **16** (2009) (1) N18.
- [13] L. Lovász, Kneser's conjecture, chromatic numbers and homotopy, *Journal of Combinatorial Theory A* **25** (1978) 319-324.
- [14] L. Lovász, *Combinatorial Problems and Exercises*, North Holland 1993.